Scientia Review, Vol. 02(03), March 2026, pp. 5-7

ISSN: 3107-7919

Scientia Review

Website: www.scientiafoundation.in



POPULAR SCIENCE ARTICLE

Marker-Assisted Selection: Unlocking Faster and Smarter Crop Improvement through DNA Science

Ragulakollu Sravanthi

Department of Genetics and Plant Breeding, Tamil Nadu Agricultural University, Coimbatore-641003

Received: 23 February 2026
Revised: 24 February 2026
Accepted: 06 March 2026
Published online: 11 March 2026

Article ID: SR01094

Citation: Sravanthi, R. (2026). Marker-Assisted Selection: Unlocking Faster and Smarter Crop Improvement Through DNA Science. *Scientia Review*, 2(3), 5-7

Abstract

Global agriculture faces unprecedented challenges due to climate change, emerging pests and diseases, shrinking arable land, and the need to feed a rapidly growing population. Conventional breeding methods, though successful, are time-consuming and often influenced by environmental factors that mask the true genetic potential of plants. Marker-Assisted Selection (MAS) has emerged as a revolutionary molecular breeding approach that integrates DNA-based markers with traditional selection methods to accelerate crop improvement. By identifying and selecting desirable genes at the seedling stage, MAS enhances precision, reduces breeding cycles, and increases selection efficiency. This technology is particularly valuable for introgressing complex traits such as disease resistance, drought tolerance, salinity tolerance, and quality attributes through gene pyramiding and backcross breeding strategies. With advances in genomics, high-throughput genotyping, and bioinformatics, MAS has significantly contributed to the development of climate-resilient, high-yielding crop varieties.

Keywords: Marker-Assisted Selection, bioinformatics, climate-resilient, phenotype, genotypes

Introduction

Marker assisted selection refers to indirect selection for a desired plant phenotype based on the presence of molecular markers linked to that trait. It is also called as Marker assisted breeding or Marker aided selection. This approach increases accuracy, efficiency, and speed in crop improvement.

Main features of MAS

- A tight linkage between molecular marker and gene of interest
- It allows heritable traits to be linked to the DNA which is responsible for controlling that trait
- Markers (AFLP, RAPD, RFLP, SSR, SNP), and differ from species to species
- The relative efficiency of MAS is greater for characters with low heritability if a large fraction of the additive genetic variance is associated with the marker loci
- It used for improvement of both oligogenic and polygenic traits.
- MAS leads to development of non-transgenic genotypes or cultivars.
- MAS permits identification of recessive alleles even in heterozygous condition and thus speed up the progress of crop improvement

Table. 1. Comparison of MAS and Phenotypic selection

Particulars	MAS	Phenotypic selection
Dissimilarities		
Markers used	Molecular markers	Morphological markers
Time required to release as a variety	very short (3-5 years)	very long (10-15 years)
Laboratory required	Sophisticated	simple
Technical skill	Very high	Medium
Accuracy		Medium to high
Cost involved		Low to medium
Similarities		
Environmental effect	No effect	Very high
Effect of gene interaction		
Mapping of QTL	Possible	Not possible
Screening of economic traits at seedling stage		

- In the MAS, RFLP markers are widely used for genetic improvement of crop plants for various economic characters.



Fig. 1. Steps in marker assisted selection



Fig. 2. Major Approaches in Marker-Assisted Selection

1. Early Generation Selection

In this approach, MAS is applied at early stages such as F_2 or F_3 generations. Plants carrying desirable genes are identified early and undesirable ones are discarded.

Advantages:

- Saves time and resources
- Fixes desirable alleles in homozygous condition
- Reduces population size for later evaluation

2. Marker-Assisted Backcrossing (MABC)

This method transfers one or a few specific genes or QTLs from a donor parent into an elite variety. There are two levels of selection:

Foreground Selection

- Identifies plants carrying the target gene
- Uses markers closely linked to the gene
- Ensures the desired gene is retained

Background Selection

- Accelerates recovery of the recurrent parent genome
- Uses markers across the genome
- Reduces number of backcross generations required

Marker-Assisted Backcrossing is especially useful for transferring resistance genes into high-performing cultivars.

3. Marker-Assisted Recurrent Selection (MARS)

MARS is used for improving complex traits controlled by multiple genes, such as grain yield, biotic stress resistance, and abiotic stress tolerance. It increases the frequency of favorable alleles in a population through repeated cycles of selection. MARS is particularly useful when multiple QTLs from different parents need to be combined. Compared to simple backcrossing, MARS is more effective for complex traits.

4. Marker-Assisted Gene Pyramiding

Gene pyramiding is the process of combining multiple resistance genes into a single genotype. This is especially important form durable disease resistance, protection against multiple pathogen races, phenotyping for multiple resistance genes at seedling stage is nearly impossible. MAS makes pyramiding feasible and precise.

Combined Approach: MAS + Phenotypic Selection

In many cases, combining molecular markers with field evaluation provides the best results.

This is useful when:

1. Some QTLs remain unidentified
2. Marker–gene linkage is not 100% accurate
3. Genotyping is cheaper than large-scale phenotyping

This integrated approach maximizes genetic gain.



Applications of MAS

1. It is very effective, efficient and rapid method of transferring resistance to biotic and abiotic stresses inn crop plants.

2. It is useful in gene pyramiding for disease and insect resistance
3. Mas being used for transfer of male sterility and photo period insensitivity into cultivated genotypes from different sources.
4. Mas also used for improvement of quality characters in different crops such as protein quality (maize), fatty acid (linoleic acid in soybean) and storage quality in vegetable and fruit crops.
5. It is more effective in introgression of desirable genes from wild into cultivated genotypes.
6. MAS has wide application for genetic improvement of oligogenic traits as compared to polygenic trait.

Limitations of MAS

Despite its advantages, MAS has certain limitations:

- Requires prior knowledge of gene–marker linkage
- Development of reliable markers can be expensive
- Limited efficiency for highly complex polygenic traits
- Recombination between marker and gene may reduce accuracy

However, advancements in genomics and high-throughput sequencing are continuously improving marker precision and reducing costs.

The Future of Marker-Assisted Breeding

- Costly technology like mas should be supported by cigar that promotes collaborative research and training on global basis. It will help in rapid spread of technology in developing countries
- MAS can be used as a tool in crop improvement programme.it can speed up the progress for development of new varieties.
- International organizations such as FAO and CGIAR should organize training programme on regional level for capacity building in MAS technology

Table. 2. Practical achievements

Crop	Disease resistance	Insect resistance	Other traits
Rice	Bacterial blight blast, rice tungro virus, gall midge	Brown plant hopper, green leaf hopper	Submergence tolerance, salt tolerant, temperature sensitive male sterility, photoperiod sensitivity, shattering resistance, grain aroma, amylose content
Maize	Northern corn blight	-	Cytoplasmic male sterility, QPM
Wheat	Leaf rust, Powdery mildew, Loose smut	Hessian fly	Cyst nematode, root lesion nematode, earliness, coleoptile colour
Sorghum	Head smut, Cyst nematode mosaic virus	-	Fertility restoration Linolenic acid
Pea	Fusarium wilt, Powdery mildew	-	Nodulation ability
Barley	Powdery mildew, stripe rust	-	Double podding