Scientia Review, Vol. 02(11), November 2026, pp. 67-68 ISSN: 3107-7919

Scientia Review

Website: www.scientiafoundation.in

POPULAR SCIENCE ARTICLE

Superfetation: exploring the limits of mammalian reproduction

A. Thangamani[1*], B. Vigneshwaran[1], A. Sabarinathan[2], V. Prabaharan[1] and A. Elango[3]
[1]Dept. of Veterinary Gynaecology and Obstetrics, Tamil Nadu Veterinary and Animal Sciences University, Tamil Nadu, India
[2]Veterinary Clinical Complex, Tamil Nadu Veterinary and Animal Sciences University, Tamil Nadu, India
[3]Veterinary College and Research institute, Tamil Nadu Veterinary and Animal Sciences University, Tamil Nadu, India
*Email: thangamtamil19@gmail.com

Received: 14 November 2026
Revised: 15 November 2026
Accepted: 15 November 2026
Published online: 16 November 2026

Article ID: SR01196

Citation: Thangamani, A., Vigneshwaran, B., Sabarinathan, A., Prabaharan, V., & Elango, A. (2026). Superfetation: exploring the limits of mammalian reproduction. *Scientia Review*, *2*(11), 67-68

Abstract

Superfetation refers to the hypothesized occurrence of a second conception while an earlier pregnancy is already established. It may result in embryos or young of different developmental ages being present in the same female. Although reports have appeared in humans, livestock, laboratory animals and wildlife, different-sized fetuses do not by themselves prove that a second conception has occurred. Delayed implantation, differences in pregnancy length, embryonic death and other reproductive conditions can produce similar observations.

Keywords: Superfetation; mammalian reproduction; embryonic diapause; superfecundation; ovulation; implantation

Introduction

Pregnancy is generally understood as a carefully coordinated process in which ovulation, fertilization, implantation and fetal development occur in sequence. Superfetation challenges this familiar pattern. The term describes a new conception taking place while an earlier pregnancy is still continuing. In theory, this can lead to two groups of developing young of different ages within the same reproductive tract. Roellig et al. (2011) in their critical review, emphasize that superfetation should be diagnosed only when a new ovulation and fertilization occur during an intact, ongoing pregnancy. This distinction is important: an unusual litter or a large difference in fetal size may have several explanations and superfetation should not be assumed simply because other explanations are not immediately obvious.

The biological conditions required for superfetation

Three major biological events must be possible at the same time:

1. **A new ovulation must occur:** The ovary has to develop and release another mature egg from the dominant follicle even though the female is already pregnant. In many mammals, the hormonal environment of pregnancy, particularly the activity of the corpus luteum and progesterone suppresses another ovarian cycle.
2. **Sperm must reach the site of fertilization:** This may require mating during pregnancy or the survival of viable sperm in the reproductive tract. The cervix, uterine environment and placenta can create physical, chemical and immune barriers to sperm movement (Harman, 1919).
3. **The uterus must permit another embryo to implant and develop:** The endometrium must remain capable of receiving a new blastocyst while supporting the existing pregnancy. The hormonal and physical conditions needed for this are not well understood.

These requirements explain why superfetation is considered unusual in mammals. It involves overcoming several coordinated reproductive safeguards, not simply mating again during pregnancy.

Phenomenon that may be misinterpreted as superfetation:

Several reproductive events can produce apparently different-aged embryos or separated births without a second conception during an active pregnancy.

1. Superfecundation

Superfecundation occurs when two or more eggs released during the same ovulation cycle are fertilized, sometimes by sperm from different males. The resulting offspring belong to the same reproductive cycle and are not evidence of a new conception during pregnancy.

2. Delayed implantation or embryonic diapause

In some mammals, a fertilized embryo pauses at

an early stage before implantation. Embryos conceived at different times may therefore implant later or a single group of embryos may remain dormant for a period (Nottle, 1976). This can alter the apparent duration of pregnancy and may make reproductive events look like superfetation.

3. Variable gestation and split parturition

Pregnancy length can vary within a species. In addition, members of one litter may sometimes be born at different times. A short interval between successive births, therefore, does not necessarily mean that a second pregnancy began before the first ended.

4. Embryonic death, resorption and abnormal pregnancy

A fetus may stop developing or die and remain in the uterus for some time before being resorbed. This can create marked differences in size among uterine contents. Conception may also occur after an earlier pregnancy has already failed; this is not superfetation in the strict sense because the first pregnancy was no longer intact.

Difficulties in diagnosing and confirming superfetation:

A convincing diagnosis requires more than observing different-sized fetuses. Investigators need reliable evidence that a second ovulation and fertilization occurred after the first pregnancy was established.

I. Ideally, they should document the reproductive history, identify separate ovulation events and their associated corpora lutea, detect embryos early and follow both groups through development.
II. Repeated ultrasound examinations and carefully timed studies can be especially useful.
III. Post-mortem examination provides only a snapshot, while reproductive events that occurred weeks earlier may be impossible to reconstruct (Kobayashi *et al*., 1996).

Conclusion

Superfetation is one of the most unusual and challenging topics in reproductive biology. It raises important questions about how the ovary, uterus, hormones and maternal immune system might permit a new conception during an existing pregnancy. Accurate timing and direct evidence of a new ovulation and fertilization during an intact pregnancy are essential. Better monitoring and modern reproductive techniques may help resolve this long-standing biological puzzle.

References

Harman, M. T. (1919). A probable case of superfetation in the cow. *The Anatomical Record, 14*, 335–336.

Kobayashi, F., Sagawa, N., Konishi, I., Tsuruta, Y., Fujiwara, H., & Mori, T. (1996). Spontaneous conception and intrauterine pregnancy in asymptomatic missed abortion of ectopic pregnancy conceived in the previous cycle. *Human Reproduction, 11*, 1347–1349.

Nottle, F. K. (1976). Superfetation in a sow and a cow. *Australian Veterinary Journal, 52*, 298.

Roellig, K., Menzies, B. R., Hildebrandt, T. B., & Goeritz, F. (2011). The concept of superfetation: a critical review on a ‘myth’ in mammalian reproduction. *Biological Reviews, 86*, 77–95.